

***Clostridium difficile* Toxins TOXIN A, TOXIN B TOXIN A AND TOXIN B TOXOIDS BINARY TOXIN**

Clostridium difficile is the causative agent of antibiotic-associated diarrhea and pseudomembranous colitis.^{1,2} It produces two major exotoxins (Toxin A and Toxin B), which are the prototypes of the family of large clostridial cytotoxins. Some strains of *C. difficile* also produce a binary ADP-ribosylating toxin (CDT) which modifies actin.³ The pathogenic role of CDT Binary Toxin in diseases induced by *C. difficile* is not clear, but about 10% of strains isolated from patients with colitis contain CDT Binary Toxin genes.^{4,5}

Clostridium difficile A and B are high molecular weight glucosyltransferases that inhibit members of the Rho family of GTPases. Toxin A, referred to as an enterotoxin, has a molecular weight of 300 kDa and an isoelectric point, pI, of 5.3. Toxin B, known as a cytotoxin, has a molecular weight of 270 kDa and a pI of 4.1.

Both Toxin A and Toxin B deactivate small GTPases such as Rho, Rac and Cdc42 by glucosylation of a threonine residue.^{6,7,8} Inhibition of these GTPases causes the shutdown of signal transduction cascades leading to depolymerization of the cytoskeleton, gene transcription of certain stress-activated protein kinases, a drop in synthesis of phosphatidylinositol 4, 5 bisphosphate and possibly even the loss of cell polarity.⁹ Loss of cytoskeletal structure results in cell rounding, and this loss of structure may account for the host's reaction to *C. difficile*.

The GTPases targeted by Toxins A and B are linked to many cell responses; however, these toxins have different physiological effects. Both toxins cause irritation to skin and increased vascular permeability.¹⁰ Only Toxin A has been shown to have a binding site on epithelial cell surface carbohydrates¹¹ which may be linked to a dramatic reaction seen in rabbit ileum.^{23,13} Toxin B is a potent cytotoxin for mammalian cells in tissue culture.^{14,15} Toxin B is at least 1,000 times more cytotoxic than Toxin A in cell rounding assays.

By inhibition of signal transduction cascades, Toxin A and Toxin B are able to block downstream responses. Toxin A has been shown to cause induction of interleukin 8.¹⁶ Toxin B may block serotonin release,¹⁷ as well as stimulate production of phospholipases C and D.^{18,19} By virtue of its ability to affect depolymerization of the cytoskeleton, Toxin B has been shown to inhibit biphasic muscle contraction.²⁰

C. difficile Toxins A and B are 63% homologous in amino acid content and have a similar three dimensional structure.²¹ The C-terminal third of each toxin is made up of sequences called clostridial repetitive oligopeptides (CROPs) which are highly antigenic. The remaining N-terminal two-thirds of Toxins A and B are less similar to each other with respect to sequence homology however, it is this portion of each protein which contains the glucosyltransferase activity.^{22, 23}

CDT Binary Toxin is a member of the ADP-ribosyltransferase family (ADPRT).³ CDT Binary Toxin belongs in the AB Binary Toxin subfamily along with C2 toxin from *C. botulinum* and Iota toxin from *C. perfringens*.²⁴ The Binary Toxin is composed of two independently produced components, the enzymatic component A subunit, CdtA (48kD), and the binding and translocation component B subunit, CdtB (94kD), which mediates cell entry of CdtA. ²⁵ The CDT Binary Toxin causes depolymerization of the actin cytoskeleton and formation of microtubule-based membrane protrusions, which are suggested to be involved in enhanced bacterial adhesion and colonization of hypervirulent *C. difficile* strains.²⁶

The CDT Binary Toxin, B subunit, CdtB, is essential for the entry of CdtA into the cytosol.²⁷ The CdtB component must be activated via cleavage, after which it can form heptamers at the cell surface and bind to specific cell-surface receptors. The cell surface receptor has been identified as lipolysis stimulated lipoprotein receptor (LSR).²⁸ Next, CdtA binds to CdtB and is taken up into the cell by receptor-mediated endocytosis. The N-terminus of CdtA is responsible for interaction with CdtB, whereas the C-terminus contains the enzymatic activity.²⁹ Based on the activity of Iota toxin, it is predicted that CDT will irreversibly ADP-ribosylate monomeric G-actin at the Arg¹⁷⁷ residue. This ADP-ribosylation will block polymerization of G-actin to F-actin and disrupt the F-actin:G-actin equilibrium which results in cell rounding and cell death.³⁰

List Biological Laboratories, Inc. provides highly purified native *C. difficile* Toxin A and Toxin B, and recombinant Binary Toxin, A and B subunits. Toxoids of Toxin A and Toxin B prepared by formaldehyde inactivation are also available.

List Labs has several antibodies generated to these *C. difficile* products including goat antisera to *C. difficile* Toxin A as well as affinity purified chicken IgY to both *C. difficile* Toxin A and Toxin B, and the *C. difficile* Binary Toxin A and B subunits. The Anti-Binary B antibody is available both with and without biotinylation.

These products are intended for research purposes and are not intended for use in humans or as diagnostic agents. For further information, please contact List Biological Laboratories, Inc.

Ordering Information

Product No.	Description	Size
152C	Toxin A from <i>C. difficile</i>	100 µg
153	Toxin A Toxoid from <i>C. difficile</i>	100 µg
154A	Toxin B Toxoid from <i>C. difficile</i>	20 µg
155A,B	Toxin B from <i>C. difficile</i>	2 µg, 20 µg
155L	Toxin B from <i>C. difficile</i> , Liquid	50 µg
157A	Binary Toxin, A Subunit	20 µg
158A	Binary Toxin, B Subunit	40 µg
752	Anti-Toxin A from <i>C. difficile</i> (Goat)	0.5 ml
753A	Anti- <i>Clostridium difficile</i> Toxin A (Chicken IgY), Liquid	0.1 mg
754A	Anti- <i>Clostridium difficile</i> Toxin B (Chicken IgY), Liquid	0.1 mg
757A	Anti- <i>Clostridium difficile</i> Binary Toxin, A Subunit (Chicken IgY), Liquid	0.1 mg
758A	Anti- <i>Clostridium difficile</i> Binary Toxin, B Subunit (Chicken IgY), Liquid	0.1 mg
759A	Biotinylated Anti- <i>Clostridium difficile</i> Binary Toxin, B Subunit (Chicken IgY), Liquid	0.1 mg

See how others have used List Labs' products on our citations page: <https://www.listlabs.com/citations>

References

1. Borriello SP. Clostridial disease of the gut. Clin. Infect. Dis. 1995; 20(Suppl. 2):242–250. [PMID:7548565](#)
2. Kelly CP, Pothoulakis C, LaMont JT. Clostridium difficile colitis. N. Engl. J. Med. 1994; 330(4):257–262. [PMID:8043060](#)
3. Popoff MR, Rubin EJ, Gill DM, Boquet P. Actin-specific ADP-ribosyltransferase produced by a Clostridium difficile strain. Infect. Immun. 1988; 56(9):2299–2306. [PMID:3137166](#)
4. Popoff M. [Molecular biology of actin-ADP-ribosylating toxins](#) in *Bacterial protein toxins*. Ed. K. Aktories and I. Just (Springer, Berlin-Heidelberg, Germany), 2000; pp 275–306.
5. Stubbs S, Rupnik M, Gibert M, Brazier J, Duerden B, Popoff M. Production of actin-specific ADP-ribosyltransferase (binary toxin) by strains of Clostridium difficile. FEMS Microbiol. Lett. 2000; 186(2):307-312. [PMID:10802189](#)

6. Aktories K, Just I. Monoglucosylation of low-molecular-mass GTP-binding Rho proteins by clostridial cytotoxins. *Trends in Cell Biol.* 1995; 5(12):441-443. [PMID:14732022](#)
7. Dillon ST, Rubin EJ, Yakubovich M, Pothoulakis C, LaMont JT, Feig LA, Gilbert RJ. Involvement of Ras-related Rho proteins in the mechanisms of action of *Clostridium difficile* toxin A and toxin B. (1995) *Infect. Immun.* 1995; 63(4):1421-1426. [PMID:7890404](#)
8. Wilkins TD, Lyerly DM. *Clostridium difficile* toxins attack Rho. *Trends in Microbiol.* 1996; 4(2):49-51. [PMID:8820565](#)
9. Mackay, D.J.G. and Hall, A. Rho GTPases. *J. Biol. Chem.* 1998; 273(33):20685-20688. [PMID:9694808](#)
10. Lyerly DM, Lockwood DE, Richardson SH, Wilkins TD. Biological activities of toxins A and B of *Clostridium difficile*. *Infect. Immun.* 1982; 35(3):1147-1150. [PMID:7068215](#)
11. Krivan HC, Clark GF, Smith DF, Wilkins TD. Cell surface binding site for *Clostridium difficile* enterotoxin: evidence for a glycoconjugate containing the sequence Gal alpha 1-3Gal beta 1-4GlcNAc. *Infect. Immun.* 1986; 53(3):573-581. [PMID:3744552](#)
12. Pothoulakis C, LaMont JT, Eglow R, Gao N, Rubins JB, Theoharides TC, Dickey BF. Characterization of rabbit ileal receptors for *Clostridium difficile* toxin A. Evidence for a receptor-coupled G protein. *J. Clin. Investig.* 1991; 88(1):119-125. [PMID:1905325](#)
13. Eglow R, Pothoulakis C, Itzkowitz S, Israel EJ, O'Keane CJ, Gong D, Gao N, Xu YL, Walker WA, LaMont JT. Diminished *Clostridium difficile* toxin A sensitivity in newborn rabbit ileum is associated with decreased toxin A receptor. *J. Clin. Investig.* 1992; 90(3):822-829. [PMID:1325998](#)
14. Aswel, JE, Ehrich M, Van Tassel RL, Tsai C, Holdeman LV, Wilkins TD. [Characterization and Comparison of *Clostridium difficile* and Other Clostridial Toxins](#) in *Microbiology*, Ed. D. Schlessinger (American Society for Microbiology, Washington, D.C.) 1979; pp 272-275.
15. Torres J, Camorlinga-Ponce M, Munoz O. Sensitivity in culture of epithelial cells from rhesus monkey kidney and human colon carcinoma to toxins A and B from *Clostridium difficile*. *Toxicon* 1992; 30(4):419-426. [PMID:1626323](#)
16. Mahida YR, Makh S, Hyde S, Gray T, Borriello SP. Effect of *Clostridium difficile* toxin A on human intestinal epithelial cells: induction of interleukin 8 production and apoptosis after cell detachment. *Gut.* 1996; 38(3):337-347. [PMID:8675084](#)
17. Prepens U, Just I, Von Eichel-Streiber C, Aktories K. Inhibition of Fc epsilon-RI-mediated activation of rat basophilic leukemia cells by *Clostridium difficile* toxin B (monoglucosyltransferase). *J. Biol. Chem.* 1996; 271(13):7324-7329. [PMID:8631752](#)
18. Schmidt M, Bienek C, Rumenapp U, Zhang C, Lummen G, Jakobs KH, Just I, Aktories K, Moos M, Von Eichel-Streiber C. A role for Rho in receptor- and G protein-stimulated phospholipase C. Reduction in phosphatidylinositol 4,5-bisphosphate by *Clostridium difficile* toxin B. *Arch. Pharm.* 1996; 354(2):87-94. [PMID:8857584](#)
19. Schmidt M, Rumenapp U, Nehls C, Ott S, Keller J, Von Eichel-Streiber C, Jacobs KH. Restoration of *Clostridium difficile* toxin-B-inhibited phospholipase D by phosphatidylinositol 4,5-bisphosphate. *Eur. J. Biochem.* 1996; 240(3):707-712. [PMID:8856074](#)
20. Lucius C, Arner A, Steusloff A, Troschka M, Hofmann F, Aktories K, Pfitzer G. *Clostridium difficile* toxin B inhibits carbachol-induced force and myosin light chain phosphorylation in guinea-pig smooth muscle: role of Rho proteins. *J. Physiol.* 1998; 506(pt 1):83-93. [PMID:9481674](#)
21. Von Eichel-Streiber C, Boquet P, Sauerborn M, Thelestam M. Large clostridial cytotoxins--a family of glycosyltransferases modifying small GTP-binding proteins. *Trends in Microbiol.* 1996; 4(10):375-382. [PMID:8899962](#)
22. Von Eichel-Streiber C, Sauerborn M, Kuramitsu HK. Evidence for a modular structure of the homologous repetitive C-terminal carbohydrate-binding sites of *Clostridium difficile* toxins and *Streptococcus mutans* glycosyltransferases. *J. Bacteriol.* 1992; 174, 6707-6710. [PMID:1307487](#)
23. Hofman F, Busch C, Prepens U, Just I, Aktories K. Localization of the glucosyltransferase activity of *Clostridium difficile* toxin B to the N-terminal part of the holotoxin. *J. Biol. Chem.* 1997; 272(17):11074-11078. [PMID:9111001](#)
24. Holbourn KP, Shone CC, Acharya KR. A family of killer toxins. Exploring the mechanism of ADP-ribosylating toxins. *FEBS J.* 2006; 273(20):4579-93. [PMID:16956368](#)

25. Perelle S., Gibert M., Bourlioux P., Corthier G., Popoff M. R. Production of a complete binary toxin (actin-specific ADP-ribosyltransferase) by *Clostridium difficile* CD196. (1997) *Infect. Immun.* 65:1402–1407. [PMID:9119480](#)
26. Schwan C, Nölke T, Kruppke AS, Schubert DM, Lang AE, Aktories K. Cholesterol- and sphingolipid-rich microdomains are essential for microtubule-based membrane protrusions induced by *Clostridium difficile* transferase (CDT). *J Biol Chem.* 2001; 286(33):29356-65. [PMID: 21705797](#)
27. Barth H, Aktories K, Popoff MR, Stiles BG. Binary bacterial toxins: biochemistry, biology, and applications of common *Clostridium* and *Bacillus* proteins. *Microbiol Mol Biol Rev.* 2004; 68(3):373-402. [PMID: 15353562](#)
28. Papatheodorou P, Carette JE, Bell GW, Schwan C, Guttenberg G, Brummelkamp TR, Aktories K. Lipolysis-stimulated lipoprotein receptor (LSR) is the host receptor for the binary toxin *Clostridium difficile* transferase (CDT). *Proc Natl Acad Sci.* 2011; 108(39):16422-7. [PMID: 21930894](#)
29. Gülke I, Pfeifer G, Liese J, Fritz M, Hofmann F, Aktories K, Barth H. Characterization of the enzymatic component of the ADP-ribosyltransferase toxin CDTa from *Clostridium difficile*. *Infect Immun.* 2001; 69(10):6004-11. [PMID: 11553537](#)
30. Davies AH, Roberts AK, Shone CC, Acharya KR. Super toxins from a super bug: structure and function of *Clostridium difficile* toxins. *Biochem J.* 2011; 436(3):517-26. [PMID: 21615333](#)